STEPS TO RESPOND TO OPIOID OVERDOSE WITH NYXOID

(Nasal spray naloxone)

STEPS TO RESPOND TO OPIOID OVERDOSE WITH NYXOID

(Nasal spray naloxone)



DANGER

Check the area is safe.



DANGER Check the area is safe.



RESPONSE Check for signs of overdose. Are they breathing?



RESPONSE Check for signs of overdose. Are they breathing?



SEND FOR HELP Call triple zero (000).

Tell them it's an overdose and you have naloxone.

It's an opioid overdose and I have naloxone.

SEND FOR HELP Call triple zero (000). Tell them it's an overdose and you have naloxone.



AIRWAY Make sure airway is open and clear.



AIRWAY Make sure airway is open and clear.




BREATHING If not breathing normally, give 2 rescue breaths.



BREATHING If not breathing normally, give 2 rescue breaths.



NALOXONE Unwrap Nyxoid device. DON'T PRESS THE BUTTON YET I



NALOXONE Unwrap Nyxoid device. DON'T PRESS THE BUTTON YET I



Insert nozzle into nostril and press button. Remove device from nostril and discard.

Insert nozzle into nostril and press button. Remove device from nostril and discard.

CPR



- Start a cycle of CPR:
- 2 breaths + 30 compressions, 5 times. Help them breathe until the Nyxoid starts working (2 to 5 minutes).



Start a cycle of CPR: 2 breaths + 30 compressions, 5 times. Help them breathe until the Nyxoid starts working (2 to 5 minutes).



Repeat step 6 and 7 using a new Nyxoid and the opposite nostril each time, until the person responds or Paramedics arrive.

It is always safe to give additional doses of Nyxoid. If you have no more Nyxoid, continue CPR.



Repeat step 6 and 7 using a new Nyxoid and the opposite nostril each time, until the person responds or Paramedics arrive.

It is always safe to give additional doses of Nyxoid. If you have no more Nyxoid, continue CPR.

AFTERCARE



Move person into recovery position. Stay with the person until Paramedics arrive. Don't leave them alone. Continue to check breathing. Repeat steps 6 and 7 if overdose returns.

AFTERCARE

CPR



Move person into recovery position. Stay with the person until Paramedics arrive. Don't leave them alone. Continue to check breathing.

Repeat steps 6 and 7 if overdose returns.



REMEMBER TO REPLACE YOUR NYXOID



Images reproduced by permission, NSW Health © 2023

Scan the QR code to find out how or visit us: PERTH: 22 / 7 Aberdeen Street BUNBURY: 97 Spencer Street SCAN ME NSW



peer based harm reduction wa

REMEMBER TO REPLACE YOUR NYXOID Scan the QR code to find out how or visit us:

PERTH: 22 / 7 Aberdeen Street Images reproduced by **BUNBURY: 97 Spencer Street** permission, NSW Health © 2023

